

A model of electroweak-scale right-handed neutrino mass

P.Q. Hung*

Dept. of Physics, University of Virginia,
382 McCormick Road, P. O. Box 400714,
Charlottesville, Virginia 22904-4714, USA

(Dated: February 2, 2008)

If neutrino masses are realized through the see-saw mechanism, can the right-handed neutrinos be produced and detected at present and future colliders? The answer is negative in the most popular see-saw scenarios for the simple reason that they are too heavy in these models. However, a simple extension of the Standard Model (SM) particle content, including mirror fermions, two $SU(2)_L$ triplet and one singlet Higgs fields, leads to a scenario in which the see-saw mechanism is realized with the Majorana mass M_R of the right-handed neutrino being of the order of the electroweak scale or smaller. A custodial $SU(2)$ symmetry arising from the two triplet Higgs fields ensures that $\rho = 1$ at tree level even when their vacuum expectation values (VEV) which determine the value of M_R , can be as large as the electroweak scale. M_R is found to obey the bound $\frac{M_Z}{2} \leq M_R < 246 \text{ GeV}$ which makes it accessible experimentally (Tevatron, LHC or ILC) since, in our scenario, ν_R 's can couple directly to the Standard Model (SM) gauge bosons.

Two of the most important experimental discoveries in the last decade are undoubtedly neutrino oscillations and the accelerating universe. Although the knowledge of individual neutrino masses is yet to be determined experimentally, the most plausible explanation for the oscillation data is the assumption that neutrinos have masses. When combined with cosmological constraints, the picture that emerges is one in which those masses are *tiny* of $O(< 1 \text{ eV})$ [1]. However, its nature- Dirac or Majorana- is unknown at the present time.

By far, the most popular scenario for neutrino masses is the celebrated see-saw mechanism [2] where neutrinos are of Majorana types. It is well-known that in this class of scenario, small neutrino masses arise because of a large hierarchy between a Dirac mass m_D (typically of the order of a charged lepton mass) which is intrinsically linked to the electroweak scale and a Majorana mass $M_R \gg m_D$, in the form m_D^2/M_R , where M_R , the Majorana mass of the “right-handed” neutrino, is typically some Grand Unified (GUT) mass scale or at least several orders of magnitude larger than the electroweak scale. The high value of M_R makes this sector inaccessible experimentally. One has to resort to indirect methods such as neutrinoless double beta decays to probe the Majorana nature of the *light* neutrino. If M_R can be found to be of the order of the electroweak scale, one could directly look for its signatures at future colliders through the production and detection of right-handed neutrinos. This possibility is realized in a model presented below.

It is not unreasonable to ask the following question: Could one obtain the see-saw mechanism strictly within the SM $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ by just extending its particle content? If it is possible to do so, what would be the constraints on the Dirac and Majorana mass scales? What would the theoretical and experimen-

tal consequences be other than providing a model for the size of the neutrino masses? It is argued in this paper that such a model can be constructed, with interesting implications: M_R cannot be larger than $\Lambda_{EW} \sim 246 \text{ GeV}$ and the Dirac mass is unrelated to Λ_{EW} . What is the most economical way to accomplish this?

The presentation will be organized as follows. It will be shown below that one can construct the see-saw mechanism by just staying within the SM *gauge group* and by simply extending the SM particle content to include mirror fermions, two additional triplet and one singlet Higgs fields. (It should be emphasized that what we mean by mirror fermions are simply fermions that behave like the SM ones but have opposite chiralities.) M_R is found to come from the VEVs of the triplet Higgs fields which can be as large as the electroweak scale because of the existence of a custodial $SU(2)$ symmetry which ensures that $\rho = 1$. The paper ends with a brief discussion of the possible signatures of our scenario such as the production of the right-handed neutrinos through the process $q + \bar{q} \rightarrow Z \rightarrow \nu_R + \nu_R$ with ν_R decays being characteristic of their Majorana nature. As we will show at the end of the paper, the characteristic signatures are like-sign dilepton events: a high energy equivalent of the neutrinoless double beta decay.

Although the model presented here stands on its own, it can be seen to fit into a grand unified model based on the group E_6 which is designed to embed the SM and a new unbroken gauge group $SU(2)_Z$ [3] as described in [4] which necessitates the introduction of the aforementioned heavy mirror fermions [5]. At this point, it is worthwhile to emphasize the fact that the main results of this paper rest on the assumption that right-handed neutrinos belong to $SU(2)_L$ doublets but not on the details of the so-called mirror fermions. In particular, the possible observation of the “light” right-handed neutrinos does not depend on the existence of heavy mirror quarks for example, although anomaly cancellation will require their existence despite the fact that they play no role in the

*pqh@virginia.edu

subsequent discussion.

Let us first start out with the following SM particle content: l_L and Φ , which are respectively the lepton doublet and the SM Higgs field. In addition, one has e_R which is $SU(2)_L$ singlet. The product $\bar{l}_L \Phi$ or $\bar{l}_L \tilde{\Phi}$ contains an $SU(2)_L$ singlet. Since e_R and ν_R are both $SU(2)_L$ singlets, the respective Yukawa couplings to the above products are allowed giving rise to the “normal” Dirac mass terms. If, however, ν_R is *not* a singlet of $SU(2)_L$ (nor a triplet for that matter), it follows that $\bar{l}_L \tilde{\Phi}$ cannot couple to ν_R . For definiteness, let us take ν_R as belonging to an $SU(2)_L$ right-handed doublet -the so-called mirror leptons- as follows:

$$l_R^M = \begin{pmatrix} \nu_R \\ e_R^M \end{pmatrix}, \quad (1)$$

where the superscript M stands for *mirror fermions*. Just as with e_R which is $SU(2)_L$ singlet, let us also assume the existence of a charged *left-handed* $SU(2)_L$ singlet mirror lepton, e_L^M (the mirror counterpart of e_R). Notice that, in this case, anomaly cancellation operates entirely within the “lepton” sector (normal and mirror) as well as within the “quark” sector. A bilinear such as $\bar{l}_L l_R^M = \bar{\nu}_L \nu_R + \bar{e}_L e_R^M$ transforms either as a singlet or as a triplet of $SU(2)_L$. Let us for the moment assume the existence of a *singlet* scalar field ϕ_S which can couple to that fermion bilinear. We have

$$\begin{aligned} \mathcal{L}_S &= g_{Sl} \bar{l}_L \phi_S l_R^M + H.c. \\ &= g_{Sl} (\bar{\nu}_L \nu_R + \bar{e}_L e_R^M) \phi_S + H.c. \end{aligned} \quad (2)$$

The SM Yukawa couplings are given by

$$\mathcal{L}_{Y1} = g_l \bar{l}_L \Phi e_R + H.c., \quad (3a)$$

$$\mathcal{L}_{Y2} = g_l^M \bar{l}_R^M \Phi e_L^M + H.c.. \quad (3b)$$

With the following VEV's:

$$\langle \phi_S \rangle = v_S, \quad (4a)$$

$$\langle \Phi \rangle = (0, v_2/\sqrt{2}), \quad (4b)$$

one obtains from Eqs. (2, 3a, 3b) the following masses and matrices

$$m_\nu^D = g_{Sl} v_S, \quad (5)$$

for the Dirac neutrino and

$$M_l = \begin{pmatrix} m_l & m_\nu^D \\ m_\nu^D & m_{lM} \end{pmatrix}, \quad (6)$$

for the charged SM and mirror leptons. The Dirac neutrino mass in (5) is obtained from (2) and, in (6), $m_l = g_l v_2/\sqrt{2}$, $m_{lM} = g_l^M v_2/\sqrt{2}$, with the off-diagonal mixing being identical to the Dirac neutrino mass from

(2). The diagonalization of (6) gives the following eigenvalues for the charged lepton and its mirror counterpart

$$\tilde{m}_l = m_l - \frac{(m_\nu^D)^2}{m_{lM} - m_l} \quad (7a)$$

$$\tilde{m}_{lM} = m_{lM} + \frac{(m_\nu^D)^2}{m_{lM} - m_l}. \quad (7b)$$

We will assume that $m_{lM} \gg m_l$. Furthermore, it will be seen below that $m_\nu^D \ll m_{lM}, m_l$ and one can easily see that the mass mixing in (6) is *negligible* giving $\tilde{m}_l \approx m_l$ and $\tilde{m}_{lM} \approx m_{lM}$.

Before turning to the subject of this paper which is neutrino masses, we wish to mention that the coupling to ϕ_S which mixes SM fermions to their mirror counterparts, also applies to the quark sector. Although the results of this paper do not depend on the existence of mirror quarks, anomaly cancellation requires it. The following short paragraph is written simply for the purpose of completeness. It is straightforward to generalize Eq. (6) to include the quarks. First, let us denote the quarks by $q_L = (u_L, d_L)$, u_R, d_R , and their mirror counterparts by $q_R^M = (u_R^M, d_R^M)$, u_L^M, d_L^M . Replacing g_{Sl} by g_{Sq} in Eq. (2) for the quark sector and noticing that $m_q = g_q v_2/\sqrt{2}$ and $m_{qM} = g_q^M v_2/\sqrt{2}$, M_q is obtained from (6) by the replacements: $m_l \rightarrow m_q$, $m_{lM} \rightarrow m_{qM}$, and $m_\nu \rightarrow m_\nu(g_{Sq}/g_{Sl})$. The eigenvalues are now $\tilde{m}_q = m_q - \frac{(m_\nu^D)^2 (g_{Sq}/g_{Sl})^2}{m_{qM} - m_q}$, $\tilde{m}_{qM} = m_{qM} + \frac{(m_\nu^D)^2 (g_{Sq}/g_{Sl})^2}{m_{qM} - m_q}$. (We will assume $m_{qM} > m_q$.) Again, one has $\tilde{m}_q \approx m_q$, $\tilde{m}_{qM} \approx m_{qM}$. One can straightforwardly generalize the above discussions to three families.

The above exercise shows that the neutrino Dirac mass can be *independent* of the electroweak scale in this simple “model”. If this were the whole story, one could simply make the Dirac mass naturally small by having a small v_S which is not constrained by any other considerations. However, the total width of the Z boson rules out that option because the addition of the right-handed neutrinos to the coupling would increase the neutrino contribution by a factor of 2. Therefore, in our scenario, the right-handed neutrinos have to be *heavier* than half the Z mass. This statement is rather general in the sense that, if ν_R transforms *non trivially* under $SU(2)_L \otimes U(1)_Y$, a small pure neutrino Dirac mass term is *forbidden*. (This is not the case if ν_R is $SU(2)_L \otimes U(1)_Y$ singlet in which case one should fine tune the Yukawa coupling, i.e. $g_\nu \sim 10^{-11}$, in order for $m_\nu \sim O(\leq eV)$.)

Since $l_R^{M,T} \sigma_2 l_R^M$ (fermion bilinear for ν_R Majorana mass term) transforms as $(1 + 3, Y/2 = -1)$ under $SU(2)_L \otimes U(1)_Y$, the appropriate Higgs field cannot be a singlet which carries a charge +1 since its VEV would break charge conservation. This leaves us with the option of a triplet Higgs field $\tilde{\chi} = (3, Y/2 = +1)$. Explicitly, $\tilde{\chi}$

is given as

$$\tilde{\chi} = \frac{1}{\sqrt{2}} \vec{\tau} \cdot \vec{\chi} = \begin{pmatrix} \frac{1}{\sqrt{2}} \chi^+ & \chi^{++} \\ \chi^0 & -\frac{1}{\sqrt{2}} \chi^+ \end{pmatrix}. \quad (8)$$

One can have a gauge invariant Yukawa coupling of the form

$$\mathcal{L}_M = g_M l_R^{M,T} \sigma_2 \tau_2 \tilde{\chi} l_R^M. \quad (9)$$

With

$$\langle \chi^0 \rangle = v_M, \quad (10)$$

one can see from (9) that the right-handed neutrino acquires a Majorana mass M_R given by

$$M_R = g_M v_M. \quad (11)$$

Notice that the use of an $SU(2)_L$ triplet had been made before in the context of the left-right symmetric model [6] and is very different from the present model. Here it is responsible for the Majorana mass of the *right-handed* neutrino.

Without further restrictions, one could also have a Yukawa coupling of the type $g_L l_L^T \sigma_2 \tau_2 \tilde{\chi} l_L$ which would give a Majorana mass $g_L v_M$ to the left-handed neutrino. Unless g_L is unnaturally fine-tuned to be very small, the presence of this term would destroy the motivation for the see-saw mechanism. To forbid its presence at tree level, one could impose the following global symmetry $U(1)_M$ under which we have

$$l_R^M, e_L^M \rightarrow e^{i\theta_M} l_R^M, e_L^M, \tilde{\chi} \rightarrow e^{-2i\theta_M} \tilde{\chi}, \phi_S \rightarrow e^{-i\theta_M} \phi_S, \quad (12)$$

with all other particles being $U(1)_M$ singlets. In consequence, this symmetry only allows the Yukawa couplings listed in (2), (3a), (3b), and (9). Furthermore, there will be no coupling of the triplet $\tilde{\chi}$ to the fermion bilinear $\bar{l}_L l_R^M$ because of $U(1)_M$. In consequence, the neutrino Dirac mass comes solely from the VEV of ϕ_S as in (2). Although this symmetry forbids the left-handed neutrino to acquire a Majorana mass at tree level, it arises at the one-loop level as given by

$$M_L = \lambda \frac{1}{16\pi^2} \frac{m_\nu^{D^2}}{M_R} \ln \frac{M_R}{M_{\phi_S}}, \quad (13)$$

where λ is the quartic coupling of ϕ_S , M_{ϕ_S} is the mass of ϕ_S and m_ν^D and M_R are given by Eq.(5) and Eq.(11) respectively. Notice that M_L as given by (13) is at most two orders of magnitude smaller than a typical see-saw light mass $m_\nu^{D^2}/M_R$ for $\lambda < 1$. To be general, we shall keep it in the mass matrix below. Another important remark is in order here. Unlike scenarios in which neutrino Majorana masses arise either from a singlet [7] or triplet Higgs field [8] and where there is an appearance of a massless Nambu-Goldstone (NG) boson- the so-called Majoron- with severe constraints, our model entails no

such NG boson as we shall see below. Also, in the aforementioned model with a Higgs triplet, a very small VEV is given to that triplet in order to maintain the approximate relationship $\rho = 1$. For a larger VEV, a custodial symmetry is required to guarantee $\rho = 1$ at tree-level, a topic to be discussed below.

The Majorana mass matrix is given by

$$\mathcal{M} = \begin{pmatrix} M_L & m_\nu^D \\ m_\nu^D & M_R \end{pmatrix}, \quad (14)$$

where m_ν^D , M_R and M_L are given by (5), (11) and (13) respectively. If $g_{SI} \sim O(g_M)$ and $v_M \gg v_S$, the eigenvalues are approximately $M_L - (m_\nu^D)^2/M_R = -(g_{SI}^2/g_M)(v_S/v_M)v_S(1-\epsilon)$ (with $\epsilon < 10^{-2}$) and M_R . What would be the consequences of the assumption $v_M \sim \Lambda_{EW}$? Before discussing the constraint from the ρ parameter which would require a custodial symmetry, let us estimate the scale v_S by using the aforementioned assumption. If $(g_{SI}^2/g_M) \sim O(1)$, the constraint $m_\nu \leq 1 \text{ eV}$ gives

$$v_S \approx \sqrt{v_M \times 1 \text{ eV}} \sim O(10^5 \text{ eV}). \quad (15)$$

In this scenario, the singlet VEV is seen to be about six orders of magnitude smaller than the electroweak scale.

One might ask about the hierarchy between the singlet VEV and the electroweak scale, namely the question about which mechanism that can exist to protect the smallness of the singlet VEV. First, let us notice that $v_S/\Lambda_{EW} \sim 10^{-6}$. This hierarchy is not as severe as the one that one encounters in a generic Grand Unified Theory where at least 13 to 14 orders of magnitude difference exists between the GUT scale and the electroweak scale. One could for instance “fine-tune” the cross coupling between the singlet and triplet Higgs fields to be less than 10^{-12} , although this may appear “unnatural”. A more interesting possibility might be a scenario in which the effective Dirac mass of the neutrino is proportional to the present value of the singlet Higgs field $\phi_S(t_0) \sim 10^5 \text{ eV}$ whose effective potential might be of the “slow-rolling” type. This type of scenario was proposed in a mass-varying neutrino model of the first reference of [9]. The true minimum might be characterized by $v_S \gg 10^5 \text{ eV}$. This possibility is under investigation.

It is well-known that an introduction of Higgs representations other than $SU(2)_L$ doublets without making sure that there is a remaining $SU(2)$ custodial symmetry would spoil the tree-level result $\rho = 1$. One of such representations is the $SU(2)_L$ triplet scalar which has been widely studied [10] for various reasons. With only one triplet, e.g. $\tilde{\chi}$, one would obtain $\rho = 2$ [11]. However, it is shown in [11] that the custodial symmetry is preserved, i.e. $\rho = 1$, if one has two triplets, one with $\tilde{\chi}(3, Y/2 = 1)$ and $\xi = (3, Y/2 = 0)$. The two triplets, when combined, form the $(3, 3)$ representation under the global $SU(2)_L \otimes SU(2)_R$ symmetry as follows

$$\chi = \begin{pmatrix} \chi^0 & \xi^+ & \chi^{++} \\ \chi^- & \xi^0 & \chi^+ \\ \chi^{--} & \xi^- & \chi^{0*} \end{pmatrix}. \quad (16)$$

The VEV of χ , namely

$$\langle \chi \rangle = \begin{pmatrix} v_M & 0 & 0 \\ 0 & v_M & 0 \\ 0 & 0 & v_M \end{pmatrix}, \quad (17)$$

breaks the global symmetry $SU(2)_L \otimes SU(2)_R$ down to the custodial $SU(2)$ and thus guaranteeing $\rho = 1$. As shown in [11], the W and Z masses have the standard expressions $M_W = g v/2$ and $M_Z = M_W/\cos\theta_W$, with $v = \sqrt{v_2^2 + 8v_M^2}$ and where $\langle \Phi \rangle = v_2/\sqrt{2}$ and $\langle \chi_0 \rangle = \langle \xi_0 \rangle = v_M$. This scenario can accommodate even the case when $v_M > v_2$. As [11] already discussed, the $U(1)_M$ symmetry in (12) is broken *explicitly* by terms in the potential which mix ξ with χ since ξ does not carry $U(1)_M$ quantum numbers. Furthermore, such an explicit breaking term is needed in order to have a proper vacuum alignment. As a result, the massless NG bosons are absent in this model. Our model also contains a singlet ϕ_S which carries a $U(1)_M$ quantum number. Unlike the “Majoron” case, our model does not generate a NG boson from the singlet since $U(1)_M$ is already explicitly broken. The discussion of the Higgs potential and its associated implications is beyond the scope of this paper and will be presented elsewhere. (As noted in [11], the Yukawa coupling (9) breaks the custodial symmetry but its contribution to ρ can be small when ν_R and e_R^M are near degenerate.) Let us notice also that there is no Majorana term similar to (9) involving ξ . The two-triplet scenario is particularly relevant to our model because it can preserve the custodial symmetry while allowing for v_M to be of the order of the electroweak scale. It is worth noticing at this point that the S parameter can be made small in models with mirror fermions and more than one Higgs doublet [4]. Also it is worth mentioning that the would-be Majoron has a mass higher than the Z boson mass and therefore does not affect the Z width.

We conclude that the right-handed neutrino mass M_R is restricted to a rather *narrow* range

$$\frac{M_Z}{2} \leq M_R < 246 \text{ GeV}, \quad (18)$$

where the lower bound comes from the experimental Z-width requirement and the upper bound is discussed above. As a result, this model is quite *predictive* in terms of “detection” of the right-handed neutrino.

The SM fermions and their mirror counterparts are contained the following E_6 representations: $\mathbf{27}_L$ and $\mathbf{27}_L^c$. The details of the organization of the SM and mirror particles in these two representations are given in [4]. For our purpose here, one needs just the parts that are relevant to the above discussions. First l_R^M , as defined in (1), which is equivalent to $l_L^{M,c}$ is grouped in $\mathbf{27}_L$ while the SM doublet l_L is put in $\mathbf{27}_L^c$. First, the neutrino Dirac mass term as written in (2) can be seen to come from $\mathbf{27}_L^{c,T} \sigma_2 \mathbf{27}_L$. Since $\mathbf{27}_L^{c,T} \sigma_2 \mathbf{27}_L \sim \mathbf{1} + \mathbf{78} + \mathbf{650}$, the above fermion bilinear can couple to a *singlet* Higgs field as follows: $\mathbf{27}_L^{c,T} \sigma_2 \mathbf{27}_L \phi_S(\mathbf{1})$,

where $\phi_S(\mathbf{1})$ is an E_6 singlet. The Majorana mass term coming from (9) can be seen to arise from $\mathbf{27}_L^T \sigma_2 \mathbf{27}_L$. Since $\mathbf{27}_L^T \sigma_2 \mathbf{27}_L \sim \mathbf{27} + \mathbf{351} + \mathbf{351}'$, one can see that the Higgs representation that contains an $SU(2)_L$ triplet which is $SU(3)_c$ and $SU(2)_Z$ singlet is $\phi(\mathbf{351}')$ with $\mathbf{351}' \supset (\mathbf{1}, \mathbf{105}')$, where $\mathbf{105}' \supset (\mathbf{1}, \mathbf{6}) \supset (\mathbf{1}, \mathbf{3})$ under $SU(3)_c \otimes SU(3)_L \otimes U(1) \rightarrow SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$.

The first immediate experimental implication of an electroweak scale M_R is the possibility of directly “detecting” the right-handed Majorana neutrino at colliders (Tevatron, LHC, or ILC). One should notice: 1) ν_R interacts with the W and Z bosons since it is part of an $SU(2)_L$ doublet; 2) both ν_R and e_R^M interact with ν_L and e_L respectively through the singlet scalar field ϕ_S . In particular, since ϕ_S is expected to have a mass of $O(10^5 \text{ eV})$, one would have the following interesting decay modes: $\nu_R \rightarrow \nu_L + \phi_S$ and $e_R^M \rightarrow e_L + \phi_S$. If the mass of ν_R is close to but less than that of e_R^M , one could have, for instance, $e_R^M \rightarrow \nu_R + e_L + \bar{\nu}_L$ followed by $\nu_R \rightarrow \nu_L + \phi_S$. Also, the heaviest ν_R could be pair-produced through $q + \bar{q} \rightarrow Z \rightarrow \nu_R + \bar{\nu}_R$ with each ν_R decaying into a lighter e_R^M plus a real or virtual W followed by $e_R^M \rightarrow e_L + \phi_S$ at a “displaced” vertex. Since ν_R is a Majorana particle, one could have e.g. $e_R^{M,-} + W^+ + e_R^{M,-} + W^+ \rightarrow e_L^- + e_L^- + W^+ + W^+ + 2\phi_S$, a *like-sign* dilepton event which is distinctively different from the Dirac case. This would constitute a high energy equivalent of the well-known neutrinoless double beta decay. The details of the phenomenology are under investigation.

One might ask whether or not the lightest ν_R might pose a problem with the total energy density. Fortunately, it is unstable because of the decay mode $\nu_R \rightarrow \nu_L + \phi_S$. ϕ_S , in turns, can annihilate each other when $T < m_{\phi_S}$ as in $\phi_S + \phi_S^* \rightarrow \nu_L + \bar{\nu}_L$. The remnants of these processes are the light ν_L 's and the sum of the masses of the latter is constrained by cosmology to be less than 1 eV. Furthermore, by the time when nucleosynthesis was supposed to take place at $T \sim \text{MeV}$, the only neutrinos that remained were the light ν_L 's whose number is restricted to be around three. The electroweak-scale right-handed neutrinos have practically disappeared by then and therefore do not affect the big bang nucleosynthesis.

There is a possibility that the decays of the triplet Higgs particles with masses close to the electroweak scale can wash out existing lepton asymmetry when $T \leq M_\chi$. If that happens then one might need a new mechanism for generating the required baryon asymmetry. Whether or not this is a problem will depend on the details of the decays of the triplet Higgs fields. This is under investigation.

Since neutrino masses, in our scenario, come from from an entirely different source ($SU(2)_L$ singlet and triplet Higgs fields instead of the SM doublet), one might expect the leptonic “CKM” matrix to be quite different from the quark CKM matrix, which appears to be the case experimentally. Since the (small) neutrino Dirac mass scale is associated with a singlet scalar, it might be tempting to

connect it with scenarios of mass-varying neutrinos [9].

In summary, a model is constructed in which the Majorana mass of the right-handed neutrinos coming from the VEVs of $SU(2)_L$ -triplet Higgs fields is found to obey $\frac{M_Z}{2} \leq M_R < 246 \text{ GeV}$, without violating the constraint $\rho = 1$ at tree level because of the presence of a custodial symmetry. Its interest lies in the possibility of producing and detecting the right-handed neutrinos at current and future colliders, a prospect that is not present with a generic see-saw scenario where M_R is typically of the

order of the GUT scale.

Acknowledgments

I would like to thank Goran Senjanovic, Vernon Barger and Paul Frampton for discussions. This work is supported in parts by the US Department of Energy under grant No. DE-A505-89ER40518.

-
- [1] See D. Spergel *et al* (WMAP3), arXiv:astro-ph/0603449, and references therein.
 - [2] P. Minkowski, Phys. Lett. B **67**, 421 (1977); M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, eds. P. van Nieuwenhuizen and D. Z. Freedman (North Holland 1979); T. Yanagida, in *Proceeding of Workshop on Unified Theory and Baryon Number in the Universe*, eds. O. Sawada and A. Sugamoto (KEK 1979); S. L. Glashow, *The future of elementary particle physics*, in *Proceedings of the 1979 Cargese Summer Institute on quarks and leptons* (M. Levy, J. -L. Basdevant, D. Speiser, J. Speiser, R. Gatsmans, and M. Jacob, eds.) Plenum Press, New York, 1980, p. 687; R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980); J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980). For recent reviews, see V. Barger, D. Marfatia, and K. Whisnant, Int. J. Mod. Phys. **E12**, 569 (2003) [arXiv:hep-ph/0308123]; R. N. Mohapatra *et al*, arXiv:hep-ph/0510213; G. Altarelli, arXiv:hep-ph/0611117, and references therein.
 - [3] P. Q. Hung, arXiv:hep-ph/0504060. P. Q. Hung, Nucl. Phys. **B747**, 55 (2006) [arXiv:hep-ph/0512282].
 - [4] P. Q. Hung and Paola Mosconi, arXiv:hep-ph/0611001.
 - [5] For a review, see J. Maalampi and M. Roos, Phys. Rept. **186**, 53 (1990).
 - [6] J. C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974); R. N. Mohapatra and J. C. Pati, *ibid.* **11**, 2558 (1975); R. N. Mohapatra and G. Sejanovic, Phys. Rev. D **12**, 1502 (1975); *ibid*, Phys. Rev. D **23**, 165 (1981), and references therein.
 - [7] Y. Chikashige, R. N. Mohapatra, and R. D. Peccei, Phys. Rev. Lett. **45**, 1926 (1980); *ibid*, Phys. Lett. B **98**, 265 (1981).
 - [8] G. B. Gelmini and R. Roncadelli, Phys. Lett. B **99**, 411 (1981). See also J. Schechter and J. W. F. Valle in [2].
 - [9] P. Q. Hung, arXiv:hep-ph/0010126; P. Gu, X. Wang, and X. Zhang, Phys. Rev. **D68**, 087301 (2003); R. Fardon, A. E. Nelson, and N. Weiner, JCAP **0410**, 005 (2004).
 - [10] H. Georgi and M. Machacek, Nucl. Phys. **B262**, 463 (1985); R. S. Chivukula and H. Georgi, Phys. Lett. B **182**, 181 (1986); P. H. Frampton, M. C. Oh, and T. Yoshikawa, Phys. Rev. **D66**, 033007 (2002) [arXiv:hep-ph/0204273]. For a recent use of a Higgs triplet, see E. Ma and U. Sarkar, Phys. Lett. B **638**, 356 (2006) [arXiv:hep-ph/0602116].
 - [11] M. S. Chanowitz and M. Golden, Phys. Lett. B **165**, 105 (1985).